

One disease, multiple symptoms

Click on the button below to explore the organ systems affected by DM1 and learn about the resulting clinical manifestations.

START HERE

One disease, multiple symptoms


Spliceopathy drives the multisystem clinical manifestations of DM1¹

Mutant *DMPK* mRNA sequesters members of the MBNL family of proteins (critical regulators of alternative splicing), leading to splicing dysregulation known as spliceopathy.¹

Dysregulated alternative splicing of pre-mRNA due to mutation of the *DMPK* gene is a molecular hallmark of DM1.¹

As MBNL proteins control the alternative splicing of genes contributing to regulation of multiple cellular pathways, individuals with DM1 experience a spectrum of clinical manifestations in various organ systems.¹

Organ systems affected by DM1²



DM1, myotonic dystrophy type 1; **DMPK**, dystrophia myotonica protein kinase; **mRNA**, messenger ribonucleic acid; **MBNL**, muscleblind-like.

References: 1. López-Martínez A, et al. *Genes (Basel)* 2020;11:1109; 2. LoRusso S, et al. *Neurotherapeutics* 2018;15:872–884; 3. Ashizawa T, et al. *Neurol Clin Pract* 2018;8:507–520; 4. Gutiérrez Gutiérrez G, et al. *Neurología* 2020;35:185–206.




One disease, multiple symptoms

The average
diagnostic delay
for adult patients
with DM1 is

6-7
YEARS¹⁻³

DM1 shows a remarkably diverse set of symptoms and wide variability of disease onset. Together, this often results in a delay in diagnosis or misdiagnosis.^{1,4}

Most common first-reported symptoms in DM1¹




The skeletal muscle and extra-skeletal muscle manifestations of DM1 have a high inter- and intra-individual variability.⁴

References:
1. Hilbert JE, et al. *J Neurol* 2013;260:2497–2504; 2. Hamel JI, et al. *Muscle Nerve* 2022;66:508–512;
3. Hagerman KA, et al. *Muscle Nerve* 2019;59:457–464; 4. Wenninger S, et al. *Front Neurol* 2018;9:303.



One disease, multiple symptoms

Tap each area to learn more about the clinical manifestations observed in DM1



CNS, central nervous system; GI, gastrointestinal.



One disease, multiple symptoms

CNS features


Up to 93% of patients with DM1 experience excessive daytime sleepiness and it is one of the most impactful symptoms on patient’s lives.^{1,2} It is often reported as the presenting symptom.³

DM1 causes changes to the brain structure due to protein deposits and mutant RNA accumulations in the nuclei of brain cells, cellular alterations including neuronal loss and gliosis, and white matter changes that result in cognitive effects.⁴

CNS features in patients with DM1



Fatigue³



Excessive daytime
sleepiness^{3,5}

PEDIATRIC



Intellectual
disability⁶



Developmental
delays⁶

ADULTS



Behavioral-
personality
changes⁴



RNA, ribonucleic acid.

References: 1. Hagerman KA, et al. *Muscle Nerve* 2019;59:457–464; 2. Hilton-Jones D. *Curr Opin Neurol* 1997;10:399–401; 3. Sansone V, et al. *J Clin Sleep Med* 2021;17:2383–2391; 4. Weijs R, et al. *Neuropathology* 2021;41:3–20; 5. Ho G, et al. *Sleep Med* 2017;32:92–96; 6. Prasad M, et al. *J Neuromuscul Dis* 2016;3:405–412.




One disease, multiple symptoms

Ocular features

Cataract formation is the most common ocular feature in DM1.¹

Cataracts in DM1 may progress faster than other cataracts leading to early-onset cataracts.²

Other ocular symptoms in adult patients with DM1¹



Reference: 1. Moshirfar M, et al. *Clin Ophthalmol* 2022;16:2837–2842;
2. MDF. Consensus-based Care Recommendations for Adults with Myotonic Dystrophy Type 1.
https://www.myotonic.org/sites/default/files/pages/files/MDF_Consensus-basedCareRecsAdultsDM1_1_21.pdf (accessed August 2024).




One disease, multiple symptoms

Endocrine features

The risk for type 2 diabetes is approximately three times higher in patients with DM1 compared with individuals without DM1.¹

Mutant *DMPK* mRNA is expressed in the testes, pituitary gland, thyroid, pancreas, and liver, and results in loss of function and abnormal processing of gene products, eg the insulin receptor.²

Endocrine abnormalities in patients with DM1²



Used with permission of The Author(s), from Winters SJ.
J Clin Endocrinol Metab. 2021;106:2819–2827;
permission conveyed through Copyright Clearance Center.

ACTH, adrenocorticotrophic hormone; **DHEAS**, dehydroepiandrosterone sulfate; **FSH**, follicle-stimulating hormone;
LH, luteinizing hormone; **mRNA**, messenger ribonucleic acid; **PTH**, parathyroid hormone.

Reference: 1. Alsaggaf R, et al. *Int J Cancer* 2020;147:785–792; 2. Winters SJ. *J Clin Endocrinol Metab* 2021;106:2819–2827.




One disease, multiple symptoms

Cardiac features

Cardiac abnormalities are the second leading cause of mortality in DM1, accounting for 20–35% of deaths.^{1–3}

Cardiac conduction disorders in DM1 are likely caused by myocyte hypertrophy, interstitial fibrosis, and fatty infiltration.⁴

Cardiac abnormalities are detected in >50% of individuals with DM1⁴



Patients with DM1 have a three-fold higher risk of sudden cardiac death than age-matched controls.⁸



References: 1. Mathieu J, et al. *Neurology* 1999;52:1658–1662; 2. de Die-Smulders CE, et al. *Brain* 1998;121:1557–1563; 3. Groh WJ, et al. *Muscle Nerve* 2011;43:648–651; 4. Petri H, et al. *Int J Cardiol* 2014;174:31–36; 5. McNally EM, Sparano D. *Heart* 2011;97:1094–1100; 6. McNally EM, et al. *J Am Heart Assoc* 2020;9:e014006; 7. Sochala M, et al. *Circulation* 2018;138:1169–1171; 8. Russo V, et al. *J Clin Med* 2023;12:1947.



One disease, multiple symptoms


Respiratory failure

Respiratory failure is the primary cause of death, accounting for mortality in 31–51% of individuals with DM1.^{1–3}

Reduced cough efficiency in DM1 is due to weakness of the inspiratory and expiratory muscles. This leads to impaired airway clearance and an increased risk of aspiration and pulmonary infections.⁴

Respiratory features of DM1

Respiratory failure typically occurs in adults between 50–60 years of age^{2,3}



Restrictive ventilatory pattern is the predominant ventilatory function abnormality in DM1⁵

Most commonly impaired respiratory parameters:^{5,6}
Forced vital capacity (FVC)
Vital capacity (VC)



References: 1. Mathieu J, et al. *Neurology* 1999;52:1658–1662; 2. de Die-Smulders CE, et al. *Brain* 1998;121:1557–1563; 3. Groh WJ, et al. *Muscle Nerve* 2011;43:648–651; 4. Boentert M, et al. *Respiration* 2020;99:360–368; 5. Hawkins AM, et al. *Neuromuscul Disord* 2019;29:198–212; 6. Hartog L, et al. *Front Neurol* 2021;12:658532.




One disease, multiple symptoms

GI features

Gastrointestinal (GI) manifestations are reported by ~80% of patients¹ with DM1 and occur along the entire digestive tract.^{2,3}

GI symptoms in DM1 are generally attributed to motility disorders caused by striated and smooth muscle dysfunction. Neurologic alterations may also play a role.^{3,4}

GI symptoms observed in DM1^{2,3}



References: 1. Heatwole C, et al. *Neurology* 2012;79:348–357; 2. Hilbert JE, et al. *Neurology* 2017;89:1348–1354; 3. Bellini M, et al. *World J Gastroenterol* 2006;12:1821–1828; 4. Peterson JAM, Cooper A. *Int J Mol Sci* 2022;23:14479.



One disease, multiple symptoms

Skeletal muscle

Please select from the options below
to learn more

Myotonia

Myopathic
face

Muscle
weakness



One disease, multiple symptoms

Skeletal muscle

Please select from the options below
to learn more

Myotonia

Myopathic
face

Muscle
weakness


88%^a of patients with DM1 experience
myotonia (difficulty relaxing muscles)¹




Parts of the body affected by myotonia²



Fingers



Tongue



Jaw

Myotonia in DM1 is thought to arise due to increased excitability of muscle fibers, leading to continuous discharges of repetitive action potentials after mechanical stimulation or voluntary contraction.²



^a Significantly higher prevalence or impact of symptoms in females (p<0.05).

References: 1. Hagerman KA, et al. *Muscle Nerve* 2019;59:457–464; 2. Wenninger S, et al. *Front Neurol* 2018;9:303; 3. Turner C, Hilton-Jones D. *J Neurol Neurosurg Psychiatry* 2010;81:358–367.



One disease, multiple symptoms

Skeletal muscle

Please select from the options below
to learn more

- Myotonia
- Myopathic face
- Muscle weakness

‘Myopathic face’ is a prominent
early feature of DM1²



Facial features in adults with DM1²



Weakness and atrophy of facial muscles
may falsely indicate a tired, sad, or emotionless patient.²

Image from Wenninger S, et al. *Front Neurol*. 2018;9:303, licensed under a CC-BY 4.0 Creative Commons license; doi: 10.3389/fneur.2018.00303.



References: 1. Hagerman KA, et al. *Muscle Nerve* 2019;59:457–464; 2. Wenninger S, et al. *Front Neurol* 2018;9:303; 3. Turner C, Hilton-Jones D. *J Neurol Neurosurg Psychiatry* 2010;81:358–367.



One disease, multiple symptoms

Skeletal muscle

Please select from the options below
to learn more

Myotonia

Myopathic
face


Muscle
weakness

Muscle weakness is reported in 94%
of adults with DM1 and is considered the
symptom with the highest impact¹



In patients with DM1, muscle biopsy demonstrates variability in
muscle fiber size, rows of internal nuclei, fibrosis, aberrant
myofibril orientation, and type I fiber atrophy.³


Distally predominant muscle weakness and atrophy are
characteristic of DM1 and mainly involve:²



Finger
flexors



Wrist
flexors



Foot
extensors



References: 1. Hagerman KA, et al. *Muscle Nerve* 2019;59:457–464; 2. Wenninger S, et al. *Front Neurol* 2018;9:303;
3. Turner C, Hilton-Jones D. *J Neurol Neurosurg Psychiatry* 2010;81:358–367.

